



1/10

FIG. 1






2/10



200710" 629E400T


3/10



4/10

FIG. 5


#	cDNA
1	1A1
2	2B1 EST
3	2B1
4	2B2
5	3A1 h.
6	3A2
7	4A1
8	β -actin
9	GAPDH
10	ext. std.



2007-10-05 10:04:59

5/10

FIG. 6



6/10

FIG. 7a





FIG. 7b



7/10

FIG. 8a




FIG. 8b




FIG. 8c




FIG. 8d




FIG. 8e






FIG. 8f



DISC-SHAPED PLATFORM

FIG. 8g

MICROTITER / NANOTITER-TYPE
 DEVICES (96/384/...)



200701062964007

200TFO" 629EH00T

8/10

PM




FIG. 9a

REGENERATION
50% UREA




FIG. 9b

MM










FIG. 9c

FIG. 10

		epi fluorescence				resonance		
	proteo-chips incubated with	image	spot means [counts]	signal/ noise*)	image	spot means [counts]	signal/ noise*)	
I)	Cy5-labelled IL6 (500ng/ml, 10 microL, 2h incubation)		46	7.0		1100	69.2	
II)	Cy5 labelled hCG (50ng/ml, 10 microL, 2h incubation)		32	5.0		646	75.1	
III)	Cy5-labelled with mixture of anti-IL6 (polyclonal) /IL6 (50ng/ml, 10 microL, 12h incubation)		25	3.8		296	44.1	

^{*}) noise [counts] defined as standard deviation of image background

Figure 11




Figure 12

